
Lee-on-Solent

([SU 552 014]–[SU 563 002])

Highlights

This site offers a range of chondrichthyan taxa comparable to that described immediately above. It has a very good range of teleost otoliths, in addition to the abundant selachian teeth and other remains.

Introduction

A diverse vertebrate fauna has been recovered from several levels within the uppermost beds of the Bracklesham Group on the lower foreshore at Lee-on-Solent. The fauna is particularly rich in shark teeth and the site has yielded over 30 species of selachians. Chimaeroids, which are poorly known elsewhere in the Middle Eocene, are represented at Lee by five species, including the type of *Elasmodus kempfi* Ward, 1976. Teleost otoliths are extremely well represented in the fossiliferous beds.

Description

The vertebrate-bearing beds occur within the uppermost part of the Bracklesham Group, the Selsey and Huntingbridge Divisions of Fisher (1862). The vertebrate fauna is composed of mammals, reptiles, birds and fishes, in association with a diverse invertebrate fauna in which crustacea and molluscs predominate.

Fauna

Marsh Farm Formation

Chondrichthyes: Elasmobranchii: Neoselachii: Squalomorphii

Heterodontus vincenti (Leriche, 1905)

H. woodwardi Casier, 1946

Isistius triturus (Winkler, 1874)

Notorhynchus kempfi Ward, 1979

Squalus minor (Leriche, 1902)

Chondrichthyes: Elasmobranchii: Neoselachii: Squatinomorphii

Squatina prima (Winkler, 1873)

Chondrichthyes: Elasmobranchii: Neoselachii: Galeomorphii

Alopias leeuensis Ward, 1978

Carcharias bopei (Agassiz, 1843)

Carcharhinus sp.

Carcharocles auriculatus (de Blainville, 1818)

Eostegostoma angusta (Nolf and Taverne *in Herman*, 1977)

Galeocerdo latidens Agassiz, 1843

Galeorhinus minor (Agassiz, 1835)

G. recticonus (Winkler, 1874)

G. cf. ypresiensis (Casier, 1946)

Hemiscyllium bruxellensis (Herman, 1977)

Isurus praecursor (Leriche, 1904)

Isurolamna affinis (Casier, 1946)

Jaeckelotodus trigonalis (Jaekel, 1895)

'*Lamna*' *lerichei* Casier, 1946

Nebrius thieliensi (Winkler, 1873)

Odontaspis winkleri (Leriche, 1905)

Physogaleus secundus (Winkler, 1874)

P. tertius (Winkler, 1874)

Scyliorhinus gilberti Casier, 1946

S. pattersoni Cappetta, 1977

Scyliorhinus spp.

'*Scyliorhinus*' *minutissimus* (Winkler, 1873)

'S.' *biauriculatus* Casier, 1950

'*Scyliorhinus*' sp.

Synodontaspis macrotus (Agassiz, 1843)

S. acutissima (Agassiz, 1843)

S. striatus (Winkler, 1874)

Chondrichthyes: Elasmobranchii: Neoselachii: Batomorphii

Aetobatus irregularis Agassiz, 1843

Aetobatus sp.

Burnhamia davesi (Woodward, 1889)

Dasyatis duponti (Winkler, 1847)

D. jaekeli (Leriche, 1905)

Dasyatis sp.

Myliobatis dixoni Agassiz, 1843

M. latidens Woodward, 1888

M. nzadinensis (Darteville and Casier, 1943)

M. striatus Buckland, 1837

M. toliapicus Agassiz, 1843

Myliobatis sp.

Pristis lathami (Galeotti, 1837) *Pristis* sp.

Propristis schweinfurthi Dames, 1883

Rhinobatos bruxellensis (Jaekel, 1894)

Rhinoptera sherbourni White, 1926

Rhynchobatus vincenti Uaekel, 1894)

Chondrichthyes: Holocephali: Chimaeriformes

Amylodon venablesae (Casier, 1966)

Callorhinchus newtoni Ward, 1973

Edaphodon bucklandi (Agassiz, 1843)

E. leptognathus Agassiz, 1843

Elasmodus kempfi Ward, 1976

Huntingbridge Division

Chondrichthyes: Elasmobranchii: Neoselachii: Squalomorphii

Heterodontus vincenti (Leriche, 1905)

H. woodwardi Casier, 1946

Notorhynchus kempfi Ward, 1979

Squalus minor (Leriche, 1902)

Isistius triturus (Winkler, 1874)

Chondrichthyes: Elasmobranchii: Neoselachii: Squatinomorphii

Squatina prima (Winkler, 1873)

Chondrichthyes: Elasmobranchii: Neoselachii: Galeomorphii

Carcharias hopei (Agassiz, 1843)

Galeocerdo latidens Agassiz, 1843

Galeorhinus minor (Agassiz, 1835)

G. recticonus (Winkler, 1874)

G. cf. ypresiensis (Casier, 1946)

Hemiscyllium bruxelliensis (Herman, 1977)

Isurus praecursor (Leriche, 1904)

Isurolamna affinis (Casier, 1946)

Jaeckelotodus trigonalis Uaekel, 1895

'*Lamna*' *lerichei* Casier, 1946

Odontaspis winkleri (Leriche, 1905)

Physogaleus secundus (Winkler, 1874)

P. tertius (Winkler, 1874)

Scyliorhinus gilberti Casier, 1946

S. pattersoni Cappetta, 1977

Scyliorhinus spp.

'*Scyliorhinus*' *minutissimus* (Winkler, 1873)

'S.' *biauriculatus* Casier, 1950

'*Scyliorhinus*' sp.

Synodontaspis macrotus (Agassiz, 1843)

S. acutissima (Agassiz, 1843)

S. striatus (Winkler, 1874)

Chondrichthyes: Elasmobranchii: Neoselachii: Batomorphii

Archaeomanta melenhorsti Herman, 1979

Aetobatus sp.

Burnhamia davesi (Woodward, 1889)

Dasyatis duponti (Winkler, 1847)

D. jaekeli (Leriche, 1905)

D. tricuspidata Casier, 1946

D. wochadunensis Ward, 1979

Myliobatis dixoni Agassiz, 1843

M. latidens Woodward, 1888

M. cf. toliapicus Agassiz, 1843

Myliobatis sp.

?*Myripristis* sp.

Pristis lathami (Galeotti, 1837)

Pristis sp.

Rhinobatos bruxelliensis (Jaekel, 1894)

Rhynchobatus vincenti Uaekel, 1894)

Chondrichthyes: Holocephali: Chimaeriformes

Edaphodon bucklandi (Agassiz, 1843)

E. leptognathus Agassiz, 1843

Bracklesham Group, undifferentiated

Indeterminate selachian dermal denticles, vertebrae, tail spines and coprolites

Osteichthyes: Actinopterygii: Acipenseroidae

Acipenser sp.

Osteichthyes: Neopterygii: Amiiformes

Amia sp.

Osteichthyes: Teleostei: Osteoglossiformes

Brychaetes sp.

Osteichthyes: Neopterygii: Pycnodontiformes

Pycnodus sp.

Osteichthyes: Acanthopterygii: Scombroidei

Cybium proosti (Storms, 1876)

Cylindracanthus rectus (Dixon, 1850)

Sphyraenodus cf. *antiquus* Casier, 1966

Trichuroides sagittidens (Winkler, 1874)

T. winkleri Casier, 1944

T. gulincki Casier, 1967

Xiphiorhynchus sp.

Osteichthyes: Acanthopterygii: Perciformes

Platylaemus colei Dixon, 1850

Platylaemus sp.

Sparcus sp.

?*Lutianus concavus* (Priem, 1912)

Osteichthyes: Elopomorpha: Anguilliformes

Conger sp.

Osteichthyes: Euteleostei: Tetradontiformes

Eotrigodon serratus (Gervais, 1852)

Interpretation

The uppermost Bracklesham Beds at Lee-on-Solent were deposited under conditions very similar to those that prevailed at Bracklesham Bay (q.v.), and the faunal list is correspondingly similar. Its richness in neoselachians and chimaeriforms is especially conspicuous and in part suggests a rich supply of food at upper and mid-water levels i.e. nekton and larger plankton.

The number of chondrichthyan species is large, with batomorphs and galeomorphs predominating, as elsewhere. Nevertheless there is an appreciable teleost element present, plus *Acipenser* and *Amia*. The scombrid teleosts and perciforms represent a fauna of active, small- to middle-sized fishes. Some genera such as *Conger*, *Acipenser* and *Amia* (which reach larger sizes) may have been somewhat migratory in habit, but it is difficult to impute such behaviour to these fossil forms.

Since most of the taxa are based on teeth, it is unlikely that taphonomic data will ever be significant. Selective sedimentation of hard parts cannot be ruled out, so that concentration of fossils may have occurred. On the other hand, there is no reliable evidence of mass mortality, nor of periodic influx of dead fish from other environments. The simplest interpretation of the fossils is that they accrued from large and flourishing fish populations.

Comparison with other localities

Comparison with the fossil yield at Bracklesham Bay (q.v.) is obvious, and similar depositional conditions and habitats are indicated. No other localities are as productive of fossil fishes at this level.

Conclusion

The conservation value of this site lies in its exceptionally rich fauna of fossil shark remains. The population of such predatory fishes would have depended upon a large mass of prey. The latter may have included other fishes as well as pelagic invertebrates and bottom-dwelling molluscs. However, the disarticulated remains and lack of taphonomic information precludes speculation about the rates and causes of mortality.

References