
Soldiers Point Bay RIGS

[GeoMôn Global Geopark original webpage](#)

RIGS Statement of Interest:

Soldiers Point Bay RIGS lies to the west of Breakwater Bay in sandstones belonging to the South Stack Group of rocks. The meta-sandstones appear to contain worm burrows similar to those reported by some workers on South Stack Moor. Their age is pertinent at the moment because new research casts doubt on their Precambrian age and a Cambrian age is now more likely. However, the South Stack Group, which was considered, by some, to be the oldest Group of Precambrian rocks on Anglesey, now appears to be the youngest and probably transposes the Precambrian/Cambrian boundary.

Geological setting/context: The Precambrian basement rocks of Anglesey and south-west LI \blacksquare n can be divided into several discrete groups, all of which were juxtaposed along a series of steep, brittle and/or ductile faults and shear zones (e.g. Dinorwic and Aber-Dinlle faults; Berw, Central Anglesey and LI \blacksquare n shear zones) collectively referred to as the Menai Strait Fault System (MSFS).

First, the Monian Supergroup consists of a thick sequence of polydeformed metasediments and meta-igneous rocks, comprising the South Stack, New Harbour and Gwna groups, the latter representing the type example of a large-scale submarine debris flow or mélangé said by some researchers to be of Lower Cambrian age. Ongoing research, however, may suggest a much older date for the Gwna Group with possible Cambrian ages being put forward for the South Stack metasediments.

Second, the Coedana Complex of central Anglesey comprises high-grade metasediments, amphibolites and gneisses, and low-grade, thermally metamorphosed hornfelses adjacent to a granite (Coedana Granite), which has recently yielded a late Precambrian zircon age of $614 \pm 4\text{Ma}$.

Third, a belt of schists and metabasites displaying blueschist facies grade of metamorphism lies within the MSFS. The metabasites exhibit a strong mid-ocean ridge basalt signature and have yielded ages of 580-590Ma.

Fourth, the Sarn Complex in LI \blacksquare n comprises metagabbros and granite rocks which occur to the south-east of the LI \blacksquare n Shear Zone (LSZ), a continuation of the MSFS, which separates these igneous rocks from low-grade Monian mélangé to the north-west

A late Precambrian zircon magmatic age of $615 \pm 2\text{Ma}$ has been obtained from a metagabbro of the LSZ. Fifth, on the mainland of north-west Wales, the Arfon Group comprises a thick sequence of tuffs and volcanoclastic rocks, dated at $614 \pm 2\text{Ma}$, which are conformably overlain by late Lower Cambrian siltstones. Correlatives of the Arfon Group may occur as isolated outliers on Anglesey and, if proven, would provide an important potential lithostratigraphical link across the MSFS. The stratigraphical correlation between the various units has proved highly controversial. The recent recognition of mylonitic rocks, for example in the LSZ, emphasises the presence of tectonic contacts and indicates that each component may represent a so-called 'suspect terrane' which was transported laterally into position along the major faults and shear zones.

Ongoing unpublished research suggests, that Anglesey's Precambrian rocks accumulated in accretionary prisms, providing a tectonic sequence rather than a stratigraphic sequence which was formerly accepted. This new research would reverse the accepted stratigraphic order of the bedded succession, since the first (oldest) material to be accreted lies above later accreted material and thus reverses the age relationships for the South Stack Group, the New Harbour Group and the Gwna Group established for the island by Robert Shackleton. This Precambrian basement later formed the north-west margin of the Lower Palaeozoic Basin, the initiation of which was contemporaneous with Arfon basement

terrane and was completed at least by early Ordovician times since an unconformable Arenig overstep sequence has been identified at several localities such as Wig Bach, Parwyd and Mountain Cottage Quarry. The Arenig sequence of Anglesey and Llŷn is considerably less deformed and metamorphosed than the underlying basement, although this distinction is not everywhere obvious.

Network context of the site: Soldiers Point Bay RIGS lies on northern side of South Stack Island adjacent to the Breakwater in Holyhead Bay. The rock is a sandstone belonging to South Stack Group of rocks which is in complete contrast to the beautifully folded schists of the New Harbour Group of rocks on the eastern side of this bay. The appearance of well-formed worm casts makes this a very important site as it provides evidence for the rocks being Cambrian in age and not Precambrian as originally thought. The bay also provides a reference section for both the South Stack and New Harbour Group of rocks and the junction between the two must be in the middle of the bay beneath the storm beach. As such this junction is also of great importance to the understanding of Precambrian evolution in Anglesey. It is possible that all Precambrian rocks in southern Britain were part of the same Avalonian subduction system as Nova Scotia, Newfoundland and Canada.

The environment in which the life forms existed. During Precambrian times, oxygen levels in the Earth's atmosphere were less than 1%, the critical level for the start of evolution on a vast scale when macrofossils entered the equation. Thus, all types of life had to exist in the sea. By Cambrian times, this critical level had been reached and there was a burst of life and a great variety of organisms evolved. Precambrian fossils were, in the main, bacteria or algae and diverse forms reflected their location, such as deep sea, associated with underwater volcanic eruptions or shallow water and intertidal conditions. The organisms could either be attached to a substrate or free-floating.

The Palaeogeography and climate. During Precambrian times, proto-Anglesey would have been part of the super continent of Pangaea, located somewhere in the region of present day Australia. Later Pangaea split into Gondwanaland and Laurasia and proto Anglesey which was attached to the ancient rocks of western North America became part of the northward-moving continent of Laurasia. The climate in Precambrian times would have been relatively cool compared to the equatorial climate experienced around 300 million years ago which was preceded and succeeded by the tropical climate experienced 100 million years either side of the Carboniferous Period. On its journey northwards it would have experienced two orogenies and one prolonged period of regional metamorphism in late Precambrian times. In addition it would, have been affected by 'Snowball Earth', a prolonged time of freezing snow and ice cover from 750 million years ago to 600 million years or at least a glacial episode in late Precambrian times. 'Global warming', caused by volcanic eruptions and the build up of methane and carbon dioxide in the atmosphere finally released the Earth from its icy grasp at the end of Precambrian times and was, in no small part, responsible for the burst of life which heralded in the Cambrian, the first period of the Palaeozoic era.

The age of the Precambrian rocks The complexity of these rocks in Anglesey and Llŷn, both in their composition and tectonic history, has led to great discrepancies in age dating. Normally, rocks are dated by the fossil content or by geochemical analysis using radioactive decay methods. Precambrian fossils are rare and can give general comparative dates. It was not until the stromatolitic sequence at Cemaes Bay was first dated that this was possible in Anglesey. Later work has found other datable fossils in the cherts of Llanddwyn and the worm casts of South Stack but, much of the work has still to be verified because different workers have produced very different dates for the same rocks. Isotope analyses have produced more accurate dates and particularly Rb/Sr and U/Pb analyses from zircons have proved useful. Collins and Buchanan have found early Cambrian ages for the South Stack Group of rocks, thought originally by Shackleton and later workers to be the oldest of the Monian deposits in the sedimentary sequence. These workers had the South Stack Group as the oldest, succeeded by the New Harbour Group and the Gwna Group as the youngest on top. Most recently, Horák (pre-publication research) has dated the limestone olistoliths in the Gwna Melange as between 650 and 700 million years old, making the Gwna the oldest Group on Anglesey, as Greenly had said in 1919. This is in agreement with Windley et. al., (in the press) who have proposed a tectonic sequence, rather than sedimentary sequence as previously proposed. This involves ocean floor sediments and rocks descending into a subduction zone and becoming accreted onto the inner continental wall. These deposits were succeeded by rocks and sediments beneath the accreted 'prisms'. Thus, unlike a sedimentary sequence where rocks are laid on top of each other, tectonically accreted rocks young down the sequence.

Soldiers Point Bay RIGS Apart from this site and the nearby South Stack RIGS, other sites demonstrating different fossil beds in different Precambrian formations are to be found on mainland Anglesey. To select RIGS to demonstrate the Precambrian evolution of Anglesey and Llŷn, three separate networks were devised. These are: 1. Precambrian stratigraphy and structures. This network includes two sub-sets: a) Precambrian sedimentary structures; and b) tectonic structures, such as folds and faults, which may have occurred during a tectonic event in Precambrian times or later, for example, during the Caledonian Orogeny; 2. Precambrian palaeontology which includes any life-form and trace fossil, such as stromatolites, sponge spicules, bacteria, worm burrows and bioturbated metasediments. Some current research suggests that some of these fossils may be Cambrian or even Ordovician in age, although other geologists dispute this. As these life-forms were previously held to be Precambrian in age, they have been included in this category; and 3. Precambrian reference sections. These aim to represent all of the important Precambrian rock types found in Anglesey and Llŷn. They include the major mapped units of Greenly (1920). The aim is to provide the best and most accessible exposure of the rock type. These can be considered as RIGS 'type sections'. Where there is a relevant metamorphic, mineralogical, sedimentary, structural or other change across an outcrop, several representative sites have been chosen. In this study, Soldiers Bay Point RIGS belongs to Network 2 (RIGS Precambrian Palaeontology; see above).

References:

- BARGHOORN, E. S. & TYLER, S. A. (1965) Microorganisms from the Gunflint Chert. *Science*, 147, 563-577.
- BLAKE, J.F. (1888) On the Monian system of rocks. *Quarterly Journal of the Geological Society of London*, 44, 271-290.
- CARNEY, J.N., HORÁK, J.M., PHARAOH, T.C., GIBBONS, W., WILSON, D., BARCLAY, W.J., BEVINS, R.E, COPE, J.C.W. & FORD, T.D. (2000) Precambrian Rocks of England and Wales. Geological Conservation Review Series No. 20. JNCC, Peterborough, 252pp.
- DOWNIE, C. C. (1974). Acritarchs from near the Precambrian — Cambrian boundary — a preliminary report. *Rev. Palaeobotany and Palynology*, 18, 57-60.
- FITCH, F. J., MILLER, J. A., & MENEISY, M. Y. (1963). Geochronological investigations on rocks from North Wales. *Nature*, London, 199, 449-451.
- GIBBONS, W. (1983). Stratigraphy, subduction and strike-slip faulting in the Mona Complex of North Wales — a review. *Proceedings of the Geologists' Association*, 94, 147-163.
- GIBBONS, W. & BALL, M. J. 1991. A discussion on Monian Supergroup stratigraphy in northwest Wales. *Journal of the Geological Society of London*, 148, 5-8.
- GIBBONS, W. & HORÁK, J. (1990). Contrasting metamorphic terranes in northwest Wales. In : D'LEMOS, R. S., STRACHAN, R. A. & TOPLEY, C. G. (eds) *The Cadomian Orogeny*. Special Publication of the Geological Society of London, 51, 315-327.
- GIBBONS, W. & MANN, A. 1983. Pre-Mesozoic lawsonite in Anglesey, northern Wales; preservation of ancient blueschists. *Geology*, 11, 3-6. GREENLY, E. (1919). The geology of Anglesey. *Memoirs of the Geological Survey of Great Britain*. HMSO, London, 980pp. (2 vols)
- GREENLY, E. (1920). 1:50,000 (and 1 inch to 1 mile) Geological Map of Anglesey. Geological Survey of Great Britain, Special Sheet No. 92 and (93 with parts of 94, 105 and 106).
- MILLER, J. A. & FITCH, F. J. (1964). Potassium-argon methods with special reference to basic igneous rocks. *Quarterly Journal of the Geological Society of London*, 120S, 55-69.
- MOORBATH, S. & SHACKLETON, R. M. (1966) Isotopic ages from the Precambrian Mona Complex of Anglesey, North Wales (Great Britain). *Earth and Planetary Science Letters*, 1, 113-117.

MUIR, M. D., BLISS, G. M. GRANT, P. R. & FISHER, M. J. (1979) Palaeontological evidence for the age of some supposedly Precambrian rocks in Anglesey, North Wales. *Journal of the Geological Society of London*. 136, 61-64.

RAST, N. (1981). Possible correlation of Precambrian rocks of Newport, Rhode Island, with those of Anglesey, Wales. *Geology*, 9, 596-601.

SHACKLETON, R. M. (1969). The Precambrian of North Wales. In WOOD, A. (ed.) *The Precambrian and Lower Palaeozoic rocks of Wales*. University of Wales Press, Cardiff, 1-22.

SHACKLETON, R. M. (1975). Precambrian rocks of Wales. In: HARRIS, A. L., SHACKLETON, R. M., WATSON, J., DOWNIE, C., HARLAND, W. B. & MOORBATH, S. (eds) *Precambrian. A correlation of Precambrian rocks in the British Isles*. Geological Society Special Report 6, 76-82.

TUCKER, R.D. & PHARAOH, T.C. (1991). U-Pb zircon ages for Late Precambrian igneous rocks in southern Britain. *Journal of the Geological Society of London*, 148, 435-43.

WOOD, D. S. (1974). Ophiolites, melanges, blueschists and ignimbrites; early Caledonian subduction in Wales? In: DOTT, R. R. & SHAVER, R. H. (eds) *Modern and Ancient Geosynclinal Sedimentation*. Society of Economic Palaeontologists and Mineralogists, Special Publication, 19, 334-344.

WOOD, M. & NICHOLLS, G. D. (1973). Precambrian stromatolite limestone from northern Anglesey. *Nature (Physical Science)*. 241, 65.

Site geometry: Site boundary